

# The Multiplicative Jordan Decomposition in Group Rings

Satya R. Arora

*Government College for Boys, Ludhiana, Punjab, India*

metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

*Center for Communications Research, 4320 Westerra Court,  
San Diego, California 92121  
E-mail: hales@ccrwest.org*

and

I. B. S. Passi

*Panjab University, Chandigarh 160014, India  
E-mail: passi@uoel.imtech.ernet.in*

*Communicated by Robert Steinberg*

Received August 8, 1997

## 1. INTRODUCTION

Let  $G$  be a finite group and  $K$  a field of characteristic 0. Then every element  $\alpha$  in the group algebra  $K[G]$  of  $G$  over  $K$  has a unique additive Jordan decomposition  $\alpha = \alpha_s + \alpha_n$  with  $\alpha_s, \alpha_n \in K[G]$ ,  $\alpha_s$  semisimple,  $\alpha_n$  nilpotent, and  $\alpha_s \alpha_n = \alpha_n \alpha_s$ . Recall that an element  $\alpha \in K[G]$  is said to be semisimple if the minimal polynomial  $m(X)$  of  $\alpha$  over  $K$  does not have repeated roots in the algebraic closure  $\bar{K}$  of  $K$ , or, equivalently, if  $m(X)$  is coprime to its (formal) derivative  $m'(X)$ . If the element  $\alpha$  is a unit in  $K[G]$ , then  $\alpha_s$  is also a unit and  $\alpha$  has a unique multiplicative Jordan decomposition  $\alpha = \alpha_s \alpha_u$ , with  $\alpha_u \in K[G]$  unipotent and  $\alpha_s \alpha_u = \alpha_u \alpha_s$ . If  $R$  is an integral domain with quotient field  $K$  and  $\alpha \in R[G]$ , then  $\alpha_s$  need not necessarily lie in  $R[G]$ . If this happens to be the case for all  $\alpha \in R[G]$  (resp. for all units in  $R[G]$ ), then we say that the *additive* (resp. *multiplicative*) *Jordan decomposition* holds in  $R[G]$ . It is easy to see that if

the AJD holds in  $R[G]$ , then so does the MJD. If the unit group  $\mathcal{U}(R[G])$  is viewed as a linear group, then the MJD is equivalent to the unit group being a splittable group [3, 13]. Finite groups  $G$  for which the AJD holds in the integral group ring  $\mathbf{Z}[G]$  have been characterized in [7, 8] and we have given some illustrative examples in [2] for the behavior of the MJD in  $\mathbf{Z}[G]$ . Our aim in this paper is to explore further the MJD property in integral group rings.

We begin by examining  $GL_n(R)$  for an arbitrary integral domain which is integrally closed and not a field. We note that  $GL_n(R)$  is splittable for  $n = 2$  and is not splittable for  $n \geq 4$ , while  $GL_3(R)$  is splittable if and only if  $R$  has the property that the difference of any two distinct units in  $R$  is again a unit in  $R$  (i.e., the units together with 0 form a subfield). This behavior naturally suggests, as in the case of the AJD, a severe restriction on the degrees of the Wedderburn components of the rational group algebra  $\mathbf{Q}[G]$  for the MJD to hold in  $\mathbf{Z}[G]$ . Indeed, we show that the degrees must all be less than or equal to 3. As a contribution toward characterizing the MJD for integral group rings, we examine several groups and show, in particular, that the MJD holds in the integral group ring  $\mathbf{Z}[Q_{4p}]$  of a generalized quaternion group  $Q_{4p}$ , and that the only dihedral groups  $D_{2n}$  of order  $2n$  for which the MJD holds in  $\mathbf{Z}[D_{2n}]$  are those with  $n = 2, 4$ , or an odd prime  $p$ .

## 2. PRELIMINARIES

The decomposition of the elements of a finite group algebra  $K[G]$ , with  $K$  a field of characteristic 0, into semisimple and nilpotent components rests on the following basic result in linear algebra.

**PROPOSITION 2.1.** *Let  $V$  be a finite-dimensional vector space over a field  $K$  of characteristic 0 and let  $\varphi \in \text{End}_K(V)$ . Then there exist unique elements  $\varphi_s$  and  $\varphi_n \in \text{End}_K(V)$  such that*

$$\varphi = \varphi_s + \varphi_n, \quad \varphi_s \varphi_n = \varphi_n \varphi_s,$$

where  $\varphi_s$  is semisimple and  $\varphi_n$  is nilpotent. Moreover,  $\varphi_s, \varphi_n$  can be expressed as polynomials in  $\varphi$  over  $K$ . If  $\varphi$  is invertible, then so is  $\varphi_s$ , and there is a unique factorization

$$\varphi = \varphi_s \varphi_u,$$

with  $\varphi_u \in \text{End}_K(V)$  unipotent and  $\varphi_s \varphi_u = \varphi_u \varphi_s$ .

An analogous result holds for square matrices over  $K$  and for elements of  $GL_n(K)$ .

For the reader's convenience, we recall a method of constructing the semisimple component  $\varphi_s$  for a given linear transformation  $\varphi \in \text{End}_K(V)$ . (See, for instance, [4], [5], or [12] for more details.) Let  $m(X) = \prod_{i=1}^r q_i^{e_i}(X)$  be the factorization in  $K[X]$  of the minimal polynomial  $m(X)$  of  $\varphi$  over  $K$  into irreducible factors. Let  $g(X) = \prod_{i=1}^r q_i(X)$ . Then  $(g(X), g'(X)) = 1$  and therefore there exist polynomials  $h(X)$  and  $k(X) \in K[X]$  such that

$$g'(X)h(X) + g(X)k(X) = 1.$$

Choose the least  $m$  such that  $2^m \geq e_i$  for  $i = 1, 2, \dots, r$ . Let  $\theta: K[X] \rightarrow K[X]$  be the  $K$ -algebra homomorphism given by

$$X \mapsto X - g(X)h(X).$$

Then  $\varphi_s = s(\varphi)$ , where  $s(X) = \theta^m(X)$ . In particular, if  $\varphi, \psi \in \text{End}_K(V)$  have the same minimal polynomial, then the polynomial  $s(X)$  is such that

$$\varphi_s = s(\varphi) \quad \text{and} \quad \psi_s = s(\psi).$$

### 3. GENERAL LINEAR GROUPS

Recall that a subgroup  $H$  of  $GL_n(K)$  is said to be *splittable* if, for every  $A \in H$ , the semisimple component  $A_s$  of  $A$  also lies in  $H$ .

Let  $R$  be an integral domain of characteristic 0 which is integrally closed in its quotient field  $K$  and  $R \neq K$ . Let  $\bar{K}$  denote the algebraic closure of  $K$ .

PROPOSITION 3.1. (i)  $GL_2(R)$  is *splittable*.

(ii)  $GL_3(R)$  is *splittable* if and only if, for every  $a, b \in \mathcal{U}(R)$ ,  $a \neq b$ ,  $a - b \in \mathcal{U}(R)$ .

(iii)  $GL_n(R)$  is *not splittable* for  $n \geq 4$ .

*Proof.* Let  $A \in GL_n(R)$  and let  $f(X)$  be the characteristic polynomial of  $A$ . If  $f(X)$  has no repeated root, then, by definition,  $A$  is semisimple. If  $f(X)$  has a root  $a$ , say, of multiplicity  $n$ , then  $a \in R$  and  $A_s = aI$ . Thus (i) follows. To prove (ii), suppose  $f(X) = (X - a)^2(X - b)$ ,  $a, b \in \bar{K}$ ,  $a \neq b$ . Then again it is easy to see that  $a, b \in \mathcal{U}(R)$ . If  $R$  has the indicated property, then

$$A_s = A - \frac{(A - aI)(A - bI)}{a - b}.$$

Consideration of the semisimple part of  $\begin{pmatrix} a & 1 & 0 \\ 0 & a & 1 \\ 0 & 0 & b \end{pmatrix}$ ,  $a, b \in \mathcal{U}(R)$ ,  $a \neq b$ , shows that condition (ii) is necessary, for the semisimple part of this matrix is  $\begin{pmatrix} a & 0 & -(a-b)^{-1} \\ 0 & a & 1 \\ 0 & 0 & b \end{pmatrix}$ . To prove (iii), let  $p \neq 0$ ,  $p \neq 0, -4$  be an element of  $R$  which is not invertible, and write  $x = p + 2$ . Consider the matrix

$$A = \begin{pmatrix} 0 & 1 & 1 & 1 \\ -1 & -x & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & -x \end{pmatrix}.$$

The minimal polynomial of  $A$  is  $(X^2 + xX + 1)^2$  and therefore

$$\begin{aligned} A_s &= A - \frac{1}{x^2 - 4}(A^2 + xA + I)(2A + xI) \\ &= \begin{pmatrix} 0 & 1 & \frac{x}{x^2 - 4} & \frac{x^2 - 2}{x^2 - 4} \\ -1 & -x & \frac{-2}{x^2 - 4} & \frac{-x}{x^2 - 4} \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & -x \end{pmatrix}, \end{aligned}$$

which does not belong to  $GL_4(R)$  since  $p = x - 2$  is not invertible in  $R$ . Finally, note that if  $A \in GL_n(R)$  and  $B = \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}$ , then  $B_s = \begin{pmatrix} A_s & 0 \\ 0 & 1 \end{pmatrix}$  and therefore if  $GL_n(R)$  is not splittable, then  $GL_r(R)$  is not splittable for all  $r \geq n$ . This completes the proof.

*Remark 3.2.* Let  $M_n(R)$  denote the ring of  $n \times n$  matrices over  $R$ . The preceding proof shows that the AJD holds in  $M_n(R)$  if  $n = 2$  and fails for  $n \geq 3$ ; that is, if  $A \in M_2(R)$ , then  $A_s \in M_2(R)$  and, for  $n \geq 3$ , there exist matrices over  $R$  whose semisimple parts do not have all their entries in  $R$ . Note that for the AJD to hold in  $M_2(R)$  one only requires that, whenever  $2\alpha$  and  $\alpha^2$  are in  $R$  for  $\alpha$  in  $K$ , then  $\alpha$  is in  $R$ —while for  $n \geq 3$ , the AJD fails for every integral domain which is not a field.

The preceding result immediately yields the following characterization for the AJD and the MJD to hold in  $R[G]$  when the ring  $R$  of coefficients is a ring  $K[X]$  of polynomials over a field  $K$  for which  $K[G]$  is a direct sum of matrix rings over fields.

**THEOREM 3.3.** *If  $G$  is a finite group and  $K$  is a field of characteristic 0 such that*

$$K[G] \simeq \bigoplus_{i=1}^h M_{n_i}(K_i),$$

*where the  $K_i$ 's are field extensions of  $K$ , then*

- (i)  $K[X][G]$  has the AJD if and only if  $n_i \leq 2$  for  $i = 1, 2, \dots, h$ ;
- (ii)  $K[X][G]$  has the MJD if and only if  $n_i \leq 3$  for  $i = 1, 2, \dots, h$ .

#### 4. IRREDUCIBLE DEGREES

**THEOREM 4.1.** *Let  $G$  be a finite group and  $R$  an integral domain of characteristic 0 which is not a field. Then, for the multiplicative Jordan decomposition to hold in  $R[G]$ , it is necessary that the degrees of the Wedderburn components of  $K[G]$ , where  $K$  is the field of fractions of  $R$ , must all be less than or equal to 3.*

*Proof.* Let  $K[G] \simeq \bigoplus_{i=1}^h M_{n_i}(D_i)$  be the Wedderburn decomposition of  $K[G]$  and assume that  $n_1 \geq 4$ . Let  $\pi_i: K[G] \rightarrow M_{n_i}(D_i)$  be the natural projections,  $i = 1, 2, \dots, h$ . Then we can choose nilpotent elements  $x, y \in R[G]$  and  $a, b \in R$  such that  $ab \neq 0$ ,

$$\pi_1(x) = \begin{pmatrix} 0 & a & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \pi_1(y) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & b & 0 \end{pmatrix},$$

$\pi_i(x) = 0 = \pi_i(y)$  for all  $i \geq 2$ . Let  $r (\neq 0) \in R$ , with  $rab \neq -4$ . Consider the elements  $u_1 = 1 + x$ ,  $u_2 = 1 + ry$ , both of which are units and therefore so is their product  $u = u_1 u_2$ . Now  $\pi_1(u) = I + A$ , where

$$A = \begin{pmatrix} 0 & a & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & 0 & rab & a \\ 0 & 0 & rb & 0 \end{pmatrix}.$$

The semisimple component of  $A$  is

$$A_s = \begin{pmatrix} 0 & 0 & 0 & a^2/rb \\ 0 & 0 & a & 0 \\ 0 & 0 & rab & a \\ 0 & 0 & rb & 0 \end{pmatrix}.$$

If  $u$  has a Jordan decomposition in  $R[G]$ , then  $\pi_1(u_s) \in \pi_1(R[G])$  and so  $a^2/rb$  lies in  $S = \pi_1(R[G])_{14}$ , the finitely generated  $R$ -submodule of  $D_1$  consisting of the  $(1, 4)$ -entries of the elements in  $\pi_1(R[G])$ . Hence  $S$  contains all of  $K$ . Let  $x_1, x_2, \dots, x_n$  be a set of generators for  $S$  over  $R$ . Choose a basis  $\{y_\lambda\}$  of  $D_1$  over  $K$  containing 1, say with  $y_1 = 1$ . Then we have  $x_i = \sum_\lambda c_{i,\lambda} y_\lambda$ , with  $c_{i,\lambda} \in K$ . For each  $k \in K \subset S$  we have  $k = \sum_i r_i x_i = \sum_i r_i \sum_\lambda c_{i,\lambda} y_\lambda = \sum_\lambda (\sum_i r_i c_{i,\lambda}) y_\lambda$ . Only the  $\lambda = 1$  term can be nonzero and we have  $k = \sum_i r_i c_{i,1}$ . Thus  $K$  is finitely generated as an  $R$ -module and hence  $K = R$ , a contradiction, and the result is proved.

Note that the latter part of the preceding argument allows us, in Theorem 3.1 of [7], to only assume that  $R$  is an integral domain which is not a field. An implicit corollary of Theorem 3.1 of [7] is that, for groups  $G$  of odd order,  $R[G]$  has the AJD if and only if  $G$  is abelian. The multiplicative analogue of this is as follows.

**COROLLARY 4.2.** *If neither 2 nor 3 divides the order of  $G$ , then  $R[G]$  has the MJD if and only if  $G$  is abelian.*

## 5. NONSPLITTABLE UNIT GROUPS

Let  $p, q$  be odd primes and let  $C_n, D_{2n}$  denote the cyclic group of order  $n$  and the dihedral group of order  $2n$ , respectively. We will determine which  $D_{2n}$  have the MJD for their integral group rings.

**PROPOSITION 5.1.** *If  $G$  is any one of the following groups, then its integral group ring does not have the multiplicative Jordan decomposition:*

- (i)  $G = \langle x, t \mid x^{pq} = t^{2^k} = 1, txt^{-1} = x^{-1} \rangle$ .
- (ii)  $G = \langle x, y, t \mid x^p = y^p = t^{2^k} = 1, xy = yx, txt^{-1} = x^{-1}, tyt^{-1} = y^{-1} \rangle$ .
- (iii)  $G = \langle x, y, t \mid x^p = y^q = t^{2^k} = 1, xy = yx, yt = ty, txt^{-1} = x^{-1} \rangle$ .
- (iv)  $G = \langle x, t \mid x^{2p} = t^{2^k} = 1, txt^{-1} = x^{-1} \rangle$ .
- (v)  $G = \langle x, t \mid x^8 = t^{2^k} = 1, txt^{-1} = x^{-1} \rangle$ .
- (vi)  $G = \langle x, t \mid x^p = t^{2^{k+1}} = 1, txt^{-1} = x^e \rangle$  where the multiplicative order of  $e \bmod p$  is 4 (hence 4 must divide  $p - 1$ ).

*In each case  $k$  is an arbitrary positive integer.*

*Proof.* We will exhibit, for each group listed in (i)–(v), a unit  $u$  in its integral group ring whose nilpotent component  $u_n$  does not have integral coefficients. The verification in each case is straightforward, and is therefore omitted. Such verification is aided by applying the standard represen-

tation of  $G$  in  $M_2(\mathbb{Z}[H])$  where  $H$  is a normal abelian subgroup of  $G$  of index 2 (see, e.g., the proof of Proposition 3.1 of [8]).

(i)  $p \neq q$ :

$$u = x^p(1 + (x^q - x^{-q})(1 + t^2 + \cdots + t^{2^k-2})(1 + t)),$$

$$u_n = \frac{1}{q}(1 + x^p + \cdots + x^{p(q-1)})(x^q - x^{-q})(1 + t^2 + \cdots + t^{2^k-2})(1 + t).$$

$p = q$ :

$$u = x^p(1 + (x - x^{-1})(1 + t^2 + \cdots + t^{2^k-2})(1 + t)),$$

$$u_n = \frac{1}{p}(1 + x^p + \cdots + x^{p(p-1)})(x - x^{-1})(1 + t^2 + \cdots + t^{2^k-2})(1 + t).$$

(ii)

$$u = y(1 + 2(x - x^{-1})(1 + t^2 + \cdots + t^{2^k-2})(1 + t)),$$

$$u_n = \frac{2}{p}(x - x^{-1})(1 + y + \cdots + y^{p-1})(1 + t^2 + \cdots + t^{2^k-2})(1 + t).$$

(iii)

$$\begin{aligned} u &= \left[ 1 + (x - x^{-1})(1 + t^2 + \cdots + t^{2^k-2})(1 + t) \right] \\ &\quad \times \left[ (1 + (x - x^{-1})(1 + t^2 + \cdots + t^{2^k-2})(1 - t) \right. \\ &\quad \left. \times (1 + y + \cdots + y^{q-1})) \right], \end{aligned}$$

$$\begin{aligned} u_n &= (x - x^{-1})(1 + t^2 + \cdots + t^{2^k-2})(1 + t) \\ &\quad \times \left( 1 - \frac{1}{q}(1 + y + \cdots + y^{q-1}) \right). \end{aligned}$$

(iv)

$$\begin{aligned} u &= \left[ 1 + (x^2 - x^{-2})(1 + t^2 + \cdots + t^{2^k-2})(1 + t) \right] \\ &\quad \times \left[ 1 - (x^2 - x^{-2})(1 - x^p)(1 + t^2 + \cdots + t^{2^k-2})(1 - t) \right], \end{aligned}$$

$$u_n = \frac{1}{2}(x^2 - x^{-2})(1 + x^p)(1 + t^2 + \cdots + t^{2^k-2})(1 + t).$$

(v)

$$u = x^2(1 + (x + 2x^3 - 2x^5 - x^7)(1 + t^2 + \cdots + t^{2^k-2})(1 + t)),$$

$$u_n = \frac{1}{2}(x - x^3 + x^5 - x^7)(1 + t^2 + \cdots + t^{2^k-2})(1 + t).$$

(vi) When  $k = 1$  the group ring  $\mathbf{Q}[G]$  has an irreducible component of degree 4 (see [8, p. 24]). The corresponding group ring for arbitrary  $k$  has the  $k = 1$  case as homomorphic image so will also have such a component. Hence Theorem 4.1 applies.

It has been shown in [2] that the MJD holds in the integral group ring of the dihedral group  $D_8$ . The  $k = 1$  case of the preceding proposition therefore immediately gives a characterization of the dihedral groups  $D_{2n}$  for which  $\mathbf{Z}[D_{2n}]$  has the MJD.

**THEOREM 5.2.** *The multiplicative Jordan decomposition holds in  $\mathbf{Z}[D_{2n}]$  if and only if  $n = 2, 4$ , or an odd prime.*

Note that Proposition 5.1, for  $k = 1$ , shows that the MJD fails for each of the groups  $D_{2pq}$ ,  $(C_p \times C_p) \rtimes C_2$  (where the generator of  $C_2$  inverts each element of  $C_p \times C_p$ ),  $C_q \times D_{2p}$ ,  $D_{4p}$ ,  $D_{16}$ , and  $C_p \rtimes C_4$  (where  $C_4$  acts faithfully on  $C_p$ ). These include all but one of the groups whose integral group rings were shown not to have the AJD in Proposition 3.1 of [8]. The one missing is the generalized quaternion group dealt with below.

## 6. GENERALIZED QUATERNION GROUPS

Again let  $p$  be an odd prime.

**THEOREM 6.1.** *The multiplicative Jordan decomposition holds in the integral group ring of  $Q_{4p} = \langle x, t \mid x^p = t^4 = 1, txt^{-1} = x^{-1} \rangle$ , the group of generalized quaternions of order  $4p$ .*

*Proof.* Consider the complex representations of  $Q_{4p}$  given by

$$\begin{aligned} \rho_1: x &\mapsto 1, & t &\mapsto 1, \\ \rho_2: x &\mapsto 1, & t &\mapsto -1, \\ \rho_3: x &\mapsto 1, & t &\mapsto i, \\ \rho_4: x &\mapsto \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix}, & t &\mapsto \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\ \rho_5: x &\mapsto \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix}, & t &\mapsto \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \end{aligned}$$

where  $i$  and  $\zeta$  are primitive fourth and  $p$ th roots of unity, respectively. From [6] we have

$$\mathbf{Q}[Q_{4p}] \simeq \mathbf{Q} \oplus \mathbf{Q} \oplus \mathbf{Q}(i) \oplus \rho_4(\mathbf{Q}[Q_{4p}]) \oplus \rho_5(\mathbf{Q}[Q_{4p}]).$$



Let

$$u = \sum_{i=0}^{p-1} \alpha_i x^i + \sum_{i=0}^{p-1} \beta_i x^i t + \sum_{i=0}^{p-1} \gamma_i x^i t^2 + \sum_{i=0}^{p-1} \delta_i x^i t^3,$$

where  $\alpha_i, \beta_i, \gamma_i, \delta_i \in \mathbf{Z}$ , be an arbitrary unit in  $\mathbf{Z}[Q_{4p}]$ . With the help of the complex representations  $\rho_i$  of  $Q_{4p}$ , one can easily show that if  $u$  is not semisimple then the nilpotent component of  $u$  is

$$u_n = \left[ \frac{1}{4} \sum_{i=1}^{(p-1)/2} ((\alpha_i + \gamma_i) - (\alpha_{p-i} + \gamma_{p-i}))(x^i - x^{-i}) \right. \\ \left. + \frac{1}{2p} \sum_{i=0}^{p-1} (p(\beta_i + \delta_i) - (\beta + \delta))x^i t \right] (1 + t^2),$$

where  $\beta = \beta_0 + \beta_1 + \cdots + \beta_{p-1}$  and  $\delta = \delta_0 + \delta_1 + \cdots + \delta_{p-1}$ . By considering the images of  $u \bmod \langle t^2 \rangle$  and  $\bmod \langle x \rangle$ , it can further be verified that  $u_n \in \mathbf{Z}[Q_{4p}]$ . This involves an application of Lemma 5 of [9] to the matrix  $\rho_4(u)$  to deduce that  $\beta_i = \delta_i$  for all  $i$ , and then working with  $\rho_5(u_n)$ .

We can also show that the integral group ring of a certain group of order 16, with presentation similar to that of  $Q_{4p}$ , has the MJD.

**THEOREM 6.2.** *The multiplicative Jordan decomposition holds in the integral group ring of  $G = \langle x, t \mid x^4 = t^4 = 1, txt^{-1} = x^{-1} \rangle$ .*

*Proof.* The group  $G$  has as homomorphic images both  $C_4 \times C_2$  (by mapping  $x^2$  to the identity) and the quaternion group  $Q_8$  (by identifying  $x^2$  and  $t^2$ ). It is known [11, Theorem 2.7] that, for each of these latter groups, every unit in the integral group ring is of the form  $+g$  or  $-g$  for some group element  $g$ . Furthermore, we can multiply any unit  $u$  in  $\mathbf{Z}[G]$  by one of the central units  $1, x^2, t^2, x^2 t^2$  without affecting its Jordan decomposability. This allows us to assume without loss of generality that (up to sign) every unit in  $\mathbf{Z}[G]$  is of the form  $u = v + (a + bx + ct + dxt)(1 - x^2)(1 + t^2)$  with  $a, b, c, d$  integers and  $v$  one of the elements  $1, x, t, xt$ . We now apply the previously mentioned standard representation of  $G$  into  $M_2(\mathbf{Z}[H])$ , where  $H = \langle x, t^2 \rangle$ , to conclude that  $u$  will be semisimple if  $v$  is either  $x, t$ , or  $xt$ . When  $v$  is 1 we also conclude that  $u$  is semisimple unless  $a = 0$  and  $b^2 = c^2 + d^2$ . Finally, when these conditions hold, we find that  $u_n = (bx + ct + dxt)(1 - x^2)(1 + t^2)$ , completing the proof since this has integral coefficients.

## 7. REMARKS

*Remark 1.* By Theorem 4.1, given a finite group  $G$ , for the MJD to hold in  $\mathbf{Z}[G]$  the degrees of the Wedderburn components of  $\mathbf{Q}[G]$  must all be less than or equal to 3. The case when the degrees are all 1 is well understood (see [1, 10]). If the degrees are all less than or equal to 2 and the Sylow 2-subgroups of  $G$  are abelian, then, proceeding as in [8], we see that, in view of Proposition 5.1,  $G$  must be of the type  $C_p \rtimes C_{2^m}$  with  $p$  an odd prime. To complete the characterization in this case, one needs to consider in the first instance such groups and the finite 2-groups.

*Remark 2.* Given a finite group  $G$ , let  $S$  (resp.  $U$ ) be the set of all semisimple (resp. unipotent) units in  $\mathbf{Z}[G]$ . Then the subgroup  $Y = \langle S, U \rangle$  generated by  $S$  and  $U$  is a normal subgroup and  $\mathcal{U}(\mathbf{Z}[G])/Y$  is a finitely generated (since  $\mathcal{U}(\mathbf{Z}[G])$  is so) and periodic group. The periodicity is a consequence of the fact that every element  $u \in \mathcal{U}(\mathbf{Z}[G])$  has a power  $u'$  with its Jordan components in  $\mathcal{U}(\mathbf{Z}[G])$ . It might be of interest to examine this quotient further.

## REFERENCES

1. S. R. Arora, Nilpotent elements in group algebras, *J. Indian Math. Soc.* **58** (1992), 99–104.
2. S. R. Arora, A. W. Hales, and I. B. S. Passi, Jordan decomposition and hypercentral units in integral group rings, *Comm. Algebra* **21** (1993), 25–35.
3. S. R. Arora and I. B. S. Passi, Central height of the unit group of an integral group ring, *Comm. Algebra* **21** (1993), 3673–3683.
4. A. Borel, “Linear Algebraic Groups,” Springer-Verlag, Berlin/New York, 1991.
5. C. Chevalley, “Théorie des Groupes de Lie II: Groupes Algébriques,” Hermann, Paris, 1951.
6. C. W. Curtis and I. Reiner “Methods of Representation Theory,” Vol. I, Wiley-Interscience, New York, 1990.
7. A. W. Hales, I. S. Luthar, and I. B. S. Passi, Partial augmentations and Jordan decomposition in group rings, *Comm. Algebra* **18** (1990), 2327–2341.
8. A. W. Hales and I. B. S. Passi, Integral group rings with Jordan decomposition, *Arch. Math.* **57** (1991), 21–27.
9. E. Kleinert, Units in  $\mathbf{Z}[Q_p]$ , *J. Number Theory* **26** (1987), 227–236.
10. S. K. Sehgal, Nilpotent elements in group rings, *Manuscripta Math.* **15** (1975), 65–80.
11. S. K. Sehgal, “Units in Integral Group Rings,” Longman, Essex, 1993.
12. M. Shirvani and B. A. F. Wehrfritz, “Skew Linear Groups,” Cambridge Univ. Press, 1986.
13. B. A. F. Wehrfritz, “Infinite Linear Groups,” Springer-Verlag, Berlin/New York, 1973.